




---

 AHELP for CIAO 3.4

## xsvmeka

Context: [sherpa](#)

*Jump to:* [Description](#) [Bugs](#) [See Also](#)

---

## Synopsis

M–G–K thermal plasma with variable abundances. XSpec model.

## Description

An emission spectrum from hot diffuse gas based on the model calculations of Mewe and Gronenschild (as amended by Kaastra). The model includes line emissions from several elements. Abundances are the number of nuclei per hydrogen nucleus relative to the Solar abundances set by the `xspecabundan` command.

### xsvmeka Parameters

Number	Name	Description
1	kT	plasma temperature in keV
2	nH	hydrogen density in $\text{cm}^{-3}$
3–16	(element)	abundances for He, C, N, O, Ne, Na, Mg, Al, Si, S, Ar, Ca, Fe, Ni with respect to Solar. Abundances are set by the <code>xspecabundan</code> command.
17	redshift	redshift, $z$
18	norm	$10^{-14} / (4 \pi (D_A(1+z))^2) \int n_e n_H dV$ , where $D_A$ is the angular size distance to the source (cm), $n_e$ is the electron density ( $\text{cm}^{-3}$ ), and $n_H$ is the hydrogen density ( $\text{cm}^{-3}$ )

This information is taken from the [XSpec User's Guide](#). Version 11.3.1 of the XSpec models is supplied with CIAO 3.2.

## Bugs

For a list of known bugs and issues with the XSPEC models, please visit the [XSPEC bugs page](#).

## See Also

*sherpa*

[atten](#), [bbody](#), [bbodyfreq](#), [beta1d](#), [beta2d](#), [box1d](#), [box2d](#), [bpl1d](#), [const1d](#), [const2d](#), [cos](#), [delta1d](#), [delta2d](#), [dered](#), [devaucouleurs](#), [edge](#), [erf](#), [erfc](#), [farf](#), [farf2d](#), [fpsf](#), [fpsf1d](#), [frmf](#), [gauss1d](#), [gauss2d](#), [gridmodel](#), [hubble](#),

